





NARMS – EB 2000

Veterinary Isolates

Fig. 6. Minimum Inhibitory Concentrations by Antimicrobial Agent for *Salmonella* Isolates from Turkey



⬇ Breakpoint

Diagnostic n=174


Slaughter n=518

NARMS – EB 2000


Veterinary Isolates

Fig. 6. Minimum Inhibitory Concentrations by Antimicrobial Agent for *Salmonella* Isolates from Turkey


Cefoxitin




Ceftiofur



Ceftriaxone



Cephalothin



⬇ Breakpoint


Diagnostic n=174

Slaughter n=518

NARMS – EB 2000

Veterinary Isolates

Fig. 6. Minimum Inhibitory Concentrations by Antimicrobial Agent for *Salmonella* Isolates from Turkey



⬇ Breakpoint


Diagnostic n=174

Slaughter n=518

NARMS – EB 2000

Veterinary Isolates

Fig. 6. Minimum Inhibitory Concentrations by Antimicrobial Agent for *Salmonella* Isolates from Turkey




⬇ Breakpoint

NARMS – EB 2000

Veterinary Isolates

Fig. 6. Minimum Inhibitory Concentrations by Antimicrobial Agent for Salmonella Isolates from Turkey



⬇ Breakpoint

Diagnostic n=174 Slaughter n=518